


☐

I'm not robot


reCAPTCHA

Continue

Duramune max 5 booster

ELANCO The Puppy Shot Booster - a subcutaneous vaccine for healthy healthy dogs 6 weeks of age or older, to promote the prevention of diseases caused by canine sopornica, infectious tooth ashepatitis, fogasa adenovirus type 2, toothy coronavirus, toothy parainfluenza, sugary parvovirus, Leptospiira canicola, Leptospiira icterohaemorrhagiae, Leptospiira grippotyphosa and Leptospiira pomona. DOSING/DOSING: Subcutaneous administration of a 1 mL dose. The recommended vaccination schedule should be started at 6 weeks of age or older. The pups should be re-vaccinated every 2-3 weeks until the age of 12 weeks. All dogs over 12 weeks of age should initially receive one dose of Duramune Max 5-CvK/4L and the second atdag 2-3 weeks later. Annual booster vaccination is recommended with one dose. DURAMURAMUNE™ MAX 5-cvK/4L A PUPPYSHOT™ BOOSTER ELANCO US CANINE DISORNIP-Adenovirus Type 2-Coronavirus-Parainfluenza-Parvovirus Vaccine Modified Live and Killed Virus Leptospiira Canicola-Grip Potyphosa-Icterohaemorrhagiae-Pomona bacterial extract For use only in animals This product has been shown to be effective in vaccinating healthy dogs 6 weeks of age or older with dog sopornitsusse, infectious dog hepatitis, Type 2 inetanovirus, canine coronavirus, canine parainfluenza, parvovirus (CPV), Leptospiira canicola, Leptospiira icterohaemorrhagiae, Leptospiira grippotyphosa and Leptospiira pomona. This product contains a CPV 2b strain that has been shown to be effective against the disease caused by CPV 2c in puppies with maternal antibody CPV. For more information on efficacy and safety data, visit productdata.aphis.usda.gov. Instructions for use • General instructions: Aseptically rehydrate Duramune Max 5 with canine coronavirus vaccine-Leptospiira bacterial extract. Subcutaneous administration of a dose of 1 mL. • Primary vaccination: The recommended vaccination schedule should be started at 6 weeks of age or older. The presence of a maternal antibody is known to interfere with the development of active immunity. Pups should be re-vaccinated every 2-3 weeks, at least until the age of 12 weeks. All dogs over 12 weeks of age should initially receive one dose of Duramune Max 5-CvK/4L and the second atdag 2-3 weeks later. The need for annual booster vaccinations for this product has not been established; consultation with a veterinarian is recommended. Duramune Max 5-CVK/4L (A Puppy Shot Booster) contains modified killed and live viruses. It vaccinates diseases caused by parvovirus, parainfluenza, coronavirus, type 2 adenovirus and dog sopornica. This vaccine is subcutaneously injected into healthy dogs at an age of 6 weeks and can even aid in the prevention of these diseases, as well as Leptospiira pomona, Leptospiira grippotyphosa, Leptospiira icterohaemorrhagiae, and Leptospiira canicola. * It's a cold boat battery. *This item cannot be transported to the 6 weeks and up. This vaccine protects against 5 harmful diseases and contains dead and live viruses. Elanco modified live and killed virus and Leptospiira Canicola-Grippotyphosa-Icterohaemorrhagiae-Pomona bacterial extract 25 x 1 mL injectable vaccine plus 25 x 1 mL vial as a dilution to inject adrenaline if an anaphylactic reaction occurs. Contact your nearest emergency pet hospital. Shake thoroughly before use. Avoid freezing. Store in the dark at 2°-7°C (35°-45°F). Write the container and all unused content. The recommended vaccination schedule should be started at 6 weeks of age or older. The presence of a maternal antibody is known to interfere with the development of active immunity. The pups should be re-vaccinated every 2-3 weeks until the age of 12 weeks. All dogs over 12 weeks of age should initially receive one dose of Duramune Max 5-CvK/4L and the second atdag 2-3 weeks later. Company: Elanco US Canine Sistemper-Adenovirus Type 2-Coronavirus-Parainfluenza-Parvovirus Vaccine Modified Live and Killed Virus Leptospiira Canicola-Grippotyphosa-Icterohaemrhagiae-Pomona Bacterial extract For use in animals Only This product has been shown to be effective in vaccinating healthy dogs 6 weeks of age or older against dog animal infectious-inhepatitis, type 2, type 2 intanovirus, dogs with coronavirus, with dogs with parainfluenza, dogs with parvovirus (CPV), Leptospiira canicola, Leptospiira icterohaemorrhagiae, Leptospiira grippotyphosa and Leptospiira pomona. The duration of immunity of the product has not been established. For more information on efficacy and safety data, visit productdata.aphis.usda.gov. This product contains a CPV 2b strain that has been shown to be effective against the disease caused by CPV 2c in puppies with maternal antibody CPV. Instructions for use • General instructions: Aseptically rehydrate Duramune Max 5 with bacterial extract Canine Coronavirus Vaccine-Leptospiira. Subcutaneous administration of a dose of 1 mL. • Primary immunisation: The presence of maternal antibodies is known to interfere with the development of active immunity in dogs and additional booster vaccinations will be required in most young animals. Initially, each dog should receive one dose and the second dose 2-3 weeks later. The need for annual booster vaccinations for this product has not been established; consultation with a veterinarian is recommended. Duramune Max 5-CvK/4L Puppyshot Booster Caution Store in the dark, between 2° and 8°C (35° to 46°F). AVOID FREEZING. LET'S SHAKE IT WELL. Do not mix it with other products unless it is on the label. This product has not been studied in pregnant animals. Deactivate all unused content before disposing. Epinephrin should be used in the event of an anaphylactic reaction. If exposed to humans, consult a doctor. Gentamicin and as a preservative. Duramune Max 5-CvK/4L The Puppyshot Puppyshot Caution In the absence of a Veterinary-Customer-Patient relationship, federal law prohibits the relabeling, repackaging, resale or redistribution of certain contents of the package. The logos of Duramune, Puppyshot, Elanco and the diagonal bar are trademarks of Elanco or its subsidiaries. © 2017 Elanco US Inc. reserves all rights. Elanco US Inc., Fort Dodge, IA 50501 Phone: 888-545-5973 VLN/PCN 196/46J9.29 25 Dose: 25 bottles vaccine plus 25 x 1 mL vial dilution YL102094 CPN: 1943008.2500 INNOVATION PATH, GREENFIELD, IN, 46140Copyright © 2020 Animalytix LLC. Updated: 2020-11-27 Duramune® Max 5-CvK/4LThe Puppyshot® Booster protects dogs from diseases caused by dog sopornica, Type 1 adenovirus (hepatitis), type 2 adenovirus (respiratory disease), parainfluenza, parvovirus and canine coronavirus, and four types of Leptospiira (L. canicola, L. icterohaemorrhagiae, L. grippotyphosa and L. pomona). Disornice, adenovirus, parainfluenza and parvovirus are modified live versions of the virus. Coronavirus and Leptospiira kill virus and killed antigens. It is recommended for the vaccination of healthy dogs aged six weeks or older. Contains twenty-five doses of 1 mL. In dogs aged six weeks or older, subcutaneous injections. Shake thoroughly before use. Package Contains25 (1 dose) vial of dry vaccine25 (1 mL) injection vaccine dilution warning Contains Gentamicin and Thimerosal as preservatives. Let's just vaccinate healthy animals. Animals incubating any disease or stress due to navigation, malnutrition, or parasitism may not achieve or maintain an adequate immune response. Do not vaccinate pregnant animals. Protection may not be fully established for all puppies less than 16 weeks of age who are free of maternal antibody interference. Epinephrin should be used in the event of an anaphylactic reaction. Storage: Store free of direct sunlight at 2-7°C. Don't freeze. Use new non-chemically sterilized needles and syringes. Use the entire content the first time you open it. Disposal: Burn the vaccine containers and all unused content. Instructions for use: Aseptically rehydrate Duramune ® with dilution solution Max 5-CvK/4L. Subcutaneous administration of a 1 mL dose. The recommended vaccination schedule should be started at six weeks of age. The presence of a maternal antibody is known to interfere with the development of active immunity. The pups should be re-vaccinated every two to three weeks until the age of 12 weeks. All dogs over 12 weeks of age should initially receive one dose of Duramune ® Max 5-CvK/4L and the second dose 2-3 weeks later. Annual booster vaccination is recommended with one dose. The Infovax-ID ® provides a simple and effective way to record information on vaccines administered to animals in veterinary practice. For vaccines requiring reconstitution, remove the both vials and both medical records of the animal. 1.) Hold the lower-right corner of the ear at the arrow labeled Peel Here between the thumb and the index finger. 2.) Pull evenly, slightly upwards, until the top of the label is parted from the vial. 3.) Place the label on the animal's medical chart. To ensure adhesion, press it down on the label. Duramune® Max 5 protects against diseases caused by dog sopornica, type 1 adenovirus (hepatitis), type 2 adenovirus (respiratory disease), parainfluenza and parvovirus. All viruses in the vaccine are live versions of the virus. It is recommended for the vaccination of healthy dogs aged six weeks or older. Contains twenty-five doses of 1 mL. In dogs aged six weeks or older, subcutaneous injections. Shake thoroughly before use. Package Contains25 (1 dose) vial of dry vaccine25 (1 mL) injection vaccine dilution warning Contains Gentamicin as a preservative. Let's just vaccinate healthy animals. Animals incubating any disease or stress due to navigation, malnutrition, or parasitism may not achieve or maintain an adequate immune response. Do not vaccinate pregnant. Protection may not be fully established for all puppies less than 16 weeks of age who are free of maternal antibody interference. Epinephrin should be used in the event of an anaphylactic reaction. Storage: Store free of direct sunlight at 2-7°C. Don't freeze. Do not mix with other vaccines. Use new non-chemically sterilized needles and syringes. Use the entire content the first time you open it. Disposal: Burn the vaccine containers and all unused content. Instructions for use: Aseptically rehydrate Duramune ® max 5 dilution solution. Subcutaneous administration of a 1 mL dose. The recommended vaccination schedule should be started at 6 weeks of age or older. The presence of a maternal antibody is known to interfere with the development of active immunity. The pups should be re-vaccinated every two to three weeks until the age of 12 weeks. Each dog over 12 weeks of age should initially take one dose of Duramune ® Max 5 and the second one 2 to 3 weeks later. Annual booster vaccination is recommended with one dose. The Infovax-ID ® provides a simple and effective way to record information on vaccines administered to animals in veterinary practice. For vaccines requiring reconstitution, remove the label from both vials and both labels must be indicated on the animal's medical table. 1.) Hold the lower-right corner of the ear at the arrow labeled Peel Here between the thumb and the index finger. 2.) Pull evenly, slightly upwards, until the top of the label is parted from the vial. 3.) Place the label on the animal's medical chart. To ensure adhesion, press it down on the label. Grip.